

KNOWLEDGE



Chemistry Topic 3 Structure and bonding


ORGANISER

Section 1: Bonding Key Terms


Ion	A charged particle formed when atoms lose or gain electrons .
Ionic bond	An electrostatic attraction between two oppositely charged ions (metal and non-metal) .
Electrostatic attraction	The attraction between a negatively charged particle and a positively charged particle.
Metals	In ionic bonding, metals lose electrons to become positively-charged ions.
Non-metals	In ionic bonding, non-metals gain electrons to become negatively-charged ions. Located on the right hand side of the periodic table.
Giant lattice	A large regular 3D structure that contains millions of bonds .
Covalent bond	A bond formed when non-metals share electrons . An electrostatic attraction between the positively charged nuclei of the bonded atoms and the electrons shared between them.
Molecule	A small group of atoms held together with covalent bonds. Not charged .
Polymer	Very large covalently bonded molecules with many repeating units .
Metallic bonding	The bonding of a metal consists of a lattice of positive ions surrounded by a sea of delocalised electrons . The metallic bond is the Electrostatic attraction between the positive ions and the delocalised electrons.
Alloy	A mixture of two or more elements, at least one of which is a metal . E.g. steel is a mixture of iron and carbon.

Section 2: Simple Covalent Molecules

Property	Reason
Low melting and boiling points (usually gases or liquids)	There are only weak intermolecular forces between the molecules which don't need much energy to overcome these forces.
Do not conduct electricity	Covalent molecules are not charged & have no free moving electrons.




Covalent dot and cross diagrams show which atoms the electrons have come from but don't show relative size of atoms or their arrangement in space.




Section 3: Giant Covalent Structures Made of Carbon

In Giant covalent compounds, all the atoms are bonded via strong covalent bonds in a giant lattice structure.



In Diamond, each **C is bonded to 4 other carbons** in a tetrahedral arrangement.



Graphite contains layers of hexagons with each carbon having 3 bonds. The extra electrons become delocalised between the layers.



Properties of Diamond

Property	Reason
Doesn't conduct electricity	Diamond doesn't contain delocalised electrons or ions .
Very hard	Each carbon bonds to 4 other carbon atoms with strong covalent bonds to form a lattice .
High melting point	A large amount of energy is needed to overcome all the strong covalent bonds in the lattice.

Properties of Graphite


Property	Reason
Conducts electricity	The delocalised electrons are free to move and carry charge through the structure.
Soft and slippery	Only weak intermolecular forces exist between layers , so layers can slide..

Section 4: Small Carbon-Based Structures



	Structure	Properties	Uses
Fullerene	Hollow-shaped , cage like structures and tubes which also contain hexagonal rings. E.g. Buckminsterfullerene (C ₆₀)	Very strong . Hollow so can contain other chemicals within it.	Drug delivery, lubricants, catalysts (large surface to volume ratio) and in electronics
Graphene	A single layer of graphite (one atom thick)	Very strong & light . Has delocalised electrons so it is able to conduct electricity .	Electronics, composites.
Carbon nanotube	Cylindrical tubes of carbon atoms that are very long compared to their diameter.	Very strong, light and flexible . Has delocalised electrons so it is able to conduct electricity .	Nanotechnology, electronics, reinforcing (e.g. tennis rackets).


Section 5: Ionic Bonding




When a metal and a non-metal react together, **the metal atom loses electrons** and becomes a **positive ion**. The **non-metal atom gains electrons** and becomes a **negative ion**. The ionic bond is a **strong electrostatic force of attraction between these oppositely charged ions**.

Property	Reason
High melting point and boiling points	Because it takes a lot of energy to overcome the many strong ionic bonds in the lattice. There is a strong electrostatic force between the positive and negative ions in the giant lattice .
Conduct electricity when liquid/ molten	Ions are able to move so there is a flow of charged ions (current).
Do not conduct electricity when solid	Ions are in fixed positions so cannot flow.


Section 6: Polymers



Section 7: Metallic Bonding



Properties of Pure Metals	
Property	Reason
High melting points	Strong electrostatic forces between the positive ions and delocalised electrons . Requires a large amount of energy to overcome.
Conduct electricity	the delocalised electrons are free to move and carry a charge .
Conduct heat	The delocalised electrons are free to move and transfer thermal energy through the structure.
Malleable	The layers are able to slide over each other so the metal can be bent and shaped. The attraction between the positive ions and delocalised electrons prevents the metal from shattering.



Section 8: Nanoparticles (triple only)

Nanoscience is the study of **small particles** that are between **1 and 100 nanometres** in size.

Nanoparticles may have properties **different** from those for the same materials in bulk because of their **high surface area to volume ratio**.

Nanoparticles may result in smaller quantities of materials e.g. catalysts being needed for industry.

Uses	Advantage
Sun cream (Zinc oxide nanoparticles)	Nanoparticles more effective at blocking suns rays. Nanoparticles are smaller than skin cells so can go through the skin into the bloodstream, Unpredictable effect on our cells?
Silver nanoparticles used in fridges, antimicrobial dressings.	Inhibit growth of microorganisms (protect against bacteria) Scientists are also worried about nanoparticles entering the environment and affecting aquatic life

Section 9: States of matter

